



Short-Lived Promise?

The Science and Policy
of Cumulative and
Short-Lived Climate
Pollutants

MYLES ALLEN

The Oxford Martin School at the University of Oxford is a world-leading centre of pioneering research that addresses global challenges. We invest in research that cuts across disciplines to address complex, global issues which cannot be understood and addressed by any one academic field alone. We support research that doesn't fit within conventional funding channels, but which could improve the wellbeing of this and future generations. We seek to make an impact through academic discovery and evidence-based policy recommendations. We aim to increase understanding of complex, global challenges through public engagement.

The Oxford Martin Programme on Resource Stewardship gathers an interdisciplinary team of philosophers, anthropologists, economists, modellers and environmental scientists to re-think how to monitor, manage, maintain and allocate globally important resources. Its aim is to work through understandings of individual and collective behaviour and current institutional practice, with a focus on how technical information is used in decision-making, to deliver a new framework for stewardship that will ensure that the world's essential resources remain available for generations to come.



Short-Lived Promise?

The Science and Policy
of Cumulative and
Short-Lived Climate
Pollutants

MYLES ALLEN

This paper reflects the views of the author, and does not necessarily reflect the position of the Oxford Martin School or the University of Oxford. Any errors or omissions are those of the author.

Immediate measures to reduce SLCP emissions could provide some climate benefit to the current generation through reduced warming over the next few decades, but would have little impact on peak warming unless CO₂ emissions are substantially reduced at the same time. Immediate reductions in CO₂ emissions would also deliver a more substantial climate benefit to future generations.

Discussions of climate policy often focus exclusively on carbon dioxide (CO₂), but these emissions are only one of the ways in which human activities affect global climate. Methane and other forms of atmospheric pollution also play an important role. International agreements, national policies and corporate strategies addressing climate change all involve setting priorities between reducing emissions of different climate pollutants.

For over 20 years, these priorities have been based on the notion of “CO₂-equivalent emissions” using a standard exchange rate or “emission metric” called the 100-year Global Warming Potential (GWP₁₀₀) to relate emissions of different greenhouse gases. Under the United Nations Framework Convention on Climate Change (UNFCCC), some countries have raised the possibility of adopting a different metric that would significantly reduce the value assigned to short-lived climate pollutants (SLCPs) like methane and thereby increase the relative emphasis on CO₂.

At the same time, initiatives such as the Climate and Clean Air Coalition are advocating an increased emphasis on SLCP reductions in climate policy. Other countries are proposing to include a broader range of SLCPs, including soot, in their UNFCCC contributions, effectively reducing the relative emphasis on CO₂. Early SLCP reductions, some proponents argue, would be cheaper and easier than reductions in CO₂. Many of the measures required to reduce SLCP emissions, such

as reducing soot emissions from biomass burning and coal-fired power plants, would also have significant co-benefits for human health and welfare, making them much easier to achieve politically. Some commentators have even argued for a more radical reframing of climate policy, away from long-term temperature goals and towards shorter-term targets for the global energy imbalance, the key driver of rates of change: this could also re-focus attention on SLCP reductions.

This is a confusing situation for non-specialists. While the overall aim of reducing greenhouse gas emissions to stabilise global climate remains the same, arguments are being made for very different policy priorities that appear to depend on a relatively obscure and technical issue: the choice of emission metric for comparing different climate pollutants. Different countries are using different metrics in determining their contributions to reducing emissions under the UNFCCC, further complicating the challenge of assessing overall progress towards the goal of stabilising temperatures.

This policy paper provides a non-technical overview of these issues, and sets out policy recommendations as countries and companies prepare for the 21st Session of the Conference of the Parties to the UNFCCC, to be held in Paris in December 2015. In doing so, the paper explains how the ‘short-lived’ versus ‘long-lived’ discussion is not really a technical issue at all, but an expression of inter-generational priorities.

Short-lived and long-lived climate pollutants

Cumulative emissions of CO₂ largely determine global mean surface warming by the late 21st century and beyond. To limit the warming they cause to 2 °C, CO₂ emissions must be limited to a cumulative budget over the entire industrial epoch of about one trillion tonnes of carbon, almost 60% of which has already been released, and net global CO₂ emissions must reach zero before global temperatures reach 2 °C.

Current emissions of both CO₂ and SLCPs such as methane and soot affect the rate and magnitude of climate change over the next few decades.

Reductions in SLCP emissions could be achieved at relatively low cost and with substantial co-benefits for agriculture and human health. The climate benefits of reduced emissions on these short timescales could, however, be comparable to natural climate variability, particularly on regional scales, and implementing SLCP reductions immediately would have little impact on peak warming unless CO₂ emissions are substantially reduced at the same time.

Emission metrics and emission trading

Any emission trading system or climate policy that addresses emissions of several different greenhouse gases together in a single ‘multi-gas basket’ requires some form of metric to specify what a given amount of one greenhouse gas is ‘worth’ in terms of another. The choice of metric to compare the impact of emissions of methane and other SLCPs with the impact of CO₂ depends on the timescale of interest. If the policy goal is to limit peak warming, it also depends on the ambition and success of future mitigation measures.

The standard GWP₁₀₀ metric provides (despite its name) an approximate indication of the relative importance of emissions of different gases to the increase in global temperatures over the next 20 to 40 years. GWP₁₀₀ is therefore a measure of impact on peak warming if and *only* if temperatures are expected to be approaching stabilisation within 40 years, for

which CO₂ emissions need to approach zero on a comparable timescale.

As long as CO₂ emissions continue to rise, the earliest possible time of peak warming remains many decades in the future and current SLCP emissions therefore have relatively little impact on peak temperatures. In this situation, policies that allow SLCP measures to be exchanged, traded or offset against CO₂ emission reductions using GWP₁₀₀ over-value the impact of SLCPs on peak warming and hence might discourage the CO₂ emission reductions that are required to stabilise temperatures. Replacing GWP₁₀₀ with a different metric would not solve this problem because any metric that is suitable for long-term impacts would be misleading for short-term impacts and vice versa. Using a metric that changes over time would help, but introduces greater complexity and uncertainty.

Recommendations: a “peak CO₂ first” strategy

Proponents of early action on SLCP emissions rightly emphasise the ‘complementary’ nature of SLCP and CO₂ mitigation, but it is important to be clear what this complementarity means: they are not two ways of achieving the same goal, but address fundamentally different goals, affecting different generations.

Early action on SLCP mitigation could affect the temperatures and climate impacts experienced by the generation of today’s decision-makers, but will have little impact on the warming experienced by future generations. Unless it is accompanied by ambitious reductions in CO₂ emissions, early SLCP mitigation will also have very little impact on eventual peak warming.

In the context of the goals of the UNFCCC, this paper makes the following recommendations:-

- Any policy to prevent dangerous anthropogenic interference in the climate system must limit cumulative emissions of the main long-lived climate pollutant, CO₂. Hence policies must aim to reduce net global CO₂ emissions to zero before temperatures reach any given limit, such as 2 °C.
- Policies reducing both CO₂ and SLCP emissions could reduce the rate and magnitude of climate change over the next few decades, but near-term SLCP reductions only affect peak warming if CO₂ emissions are reduced at the same time.

- No single metric can represent both the short-term impact of SLCP emissions and the cumulative impact of CO₂ emissions. The standard GWP₁₀₀ measures impact on peak warming if and only if net global CO₂ emissions are expected to be approaching zero within the next few decades.
- Rather than adjusting metrics in a single ‘multi-gas basket’ framework, specific policies are required to ensure that global CO₂ emissions are contained within a cumulative budget consistent with meeting the 2 °C goal. These policies must be independent of, and in addition to, any multi-gas emission goals.
- In effect, this implies a ‘peak CO₂ first’ strategy: the need to limit cumulative CO₂ emissions would over-ride most opportunities to offset CO₂ reductions against SLCP reduction measures until global CO₂ emissions are falling.
- As soon as CO₂ emissions are falling fast enough that there is a realistic prospect of meeting the cumulative budget, SLCP emission reductions will become a crucial priority to limit peak warming.

Introduction

The latest Scientific Assessment Report of the Intergovernmental Panel on Climate Change (IPCC) has re-stated that substantial reductions in greenhouse gas emissions will be required to meet the internationally agreed goal of avoiding more than 2 °C of global warming relative to pre-industrial temperatures.¹ Yet emissions of the principal greenhouse gas, carbon dioxide (CO₂), continue to rise.²

The challenges of securing international agreement, the perceived cost of CO₂ emission reductions, and the recognition that even relatively ambitious CO₂ measures may take decades to have a substantial impact on rising global temperatures, have together encouraged renewed interest³ in reducing emissions of other agents that cause global warming, so-called short-lived climate pollutants (SLCPs), including methane, black carbon (soot) and tropospheric ozone.^{4,5}

Some commentators have also begun to question an exclusive focus on an “effectively unachievable” global temperature target⁶ and propose replacing the 2 °C goal with targets for global energy imbalance in 2030 or 2050. One consequence of such a change of policy focus could be an increased emphasis on SLCP mitigation at the expense of CO₂.

A key attraction of action on SLCPs is that many of the measures required to reduce emissions of these agents are relatively low cost (such as reducing methane leakage from fossil fuel extraction and transport) and offer substantial co-benefits (such as improved health in developing countries from reduced soot emissions), making them potentially attractive even to governments unwilling to commit to stringent economy-wide climate targets.⁷ All countries include methane in their contributions to the United Nations Framework Convention on Climate Change (UNFCCC), and some have begun to include black carbon.⁸

At the same time, the IPCC also emphasises⁹ that, in the long-term, global temperatures are overwhelmingly determined by cumulative emissions

of CO₂ over the entire ‘Anthropocene’ epoch, not the rate of emission of greenhouse gases in any given decade.^{10,11} This recognition of the long-term importance of CO₂ has prompted renewed interest in the question of whether methane emissions, in particular, may be ‘over-valued’ in current climate policies, including the emission trading systems and other measures established by the UNFCCC.^{12,13}

For the policy community, these appear to be two contradictory developments. On the one hand, the argument is being made that climate policies should place more emphasis on reducing emissions of SLCPs like methane. On the other hand, a counter-argument is made that methane emissions are already over-valued in the existing climate policy framework and that the main focus of efforts should be on CO₂.

This policy paper sets out to explain this apparently paradoxical situation. In resolving the paradox, this paper will show how reducing SLCP emissions and reducing CO₂ emissions are not alternative ways of achieving the same goal, but should rather be seen as achieving different goals, both of which are desirable.

It is important that the significant opportunities for reducing SLCP emissions are grasped, particularly as the emerging economies from which many of these emissions originate play an increasingly prominent role in the international climate regime. But it is also important that these opportunities do not undermine progress in reducing and ultimately eliminating CO₂ emissions, which is essential in order to avoid dangerous climate change in the longer term.

To meet both of these objectives, mitigation policies must recognise the unique importance of *cumulative* carbon emissions.¹⁴ This limits the degree to which CO₂ emission reductions can be exchanged or traded for SLCP emission reductions until CO₂ emissions are falling fast enough that there is a realistic prospect of the cumulative budget being met, and hence a realistic estimate of the remaining time to peak warming.

1. Defining long-lived climate pollutants, short-lived climate pollutants and emission metrics

1.1 The main long-lived climate pollutants

Carbon dioxide (CO₂) is the single most important greenhouse gas directly affected by human activity. About 38 billion tonnes of CO₂ were emitted in 2011, over 90% from the combustion of fossil fuels and cement production, the remainder from deforestation.¹⁵ Cumulative CO₂ emissions since 1750 amount to 2,000 billion tonnes. About 45% remains in the atmosphere, raising concentrations from 278ppm in 1750 to over 390ppm in 2011. CO₂ has a very long atmospheric lifetime: 15–40% of the increase in CO₂ due to fossil fuel consumption is expected to remain in the atmosphere for over 1,000 years.¹⁶ Hence there is no sustainable CO₂ emission level: global temperatures will continue to rise until net CO₂ emissions are reduced close to zero, with peak temperatures largely determined by cumulative

CO₂ emissions up to that time. Reducing CO₂ emissions will require substantial and sustained changes to the global energy system.¹⁷

Nitrous oxide (N₂O) is the next most important of the long-lived climate pollutants, with a lifetime of about 120 years.¹⁸ N₂O is mainly generated from artificial fertiliser use in agriculture. Tonne for tonne, N₂O is a more powerful greenhouse gas than CO₂, but emissions are much lower so the stock of N₂O in the atmosphere has much less impact on global climate than CO₂ at present. Substantially reducing N₂O emissions below today's levels is expected to be challenging because of their role in food production.¹⁹

1.2 The main short-lived climate pollutants

Methane (CH₄), the main constituent of natural gas, is the most important greenhouse gas directly affected by human activity after CO₂. Methane is released by a range of activities: over the past decade, 40% came from agricultural sources such as rice paddies and livestock, 30% from fossil fuel production and use (in particular the release of natural gas), 20% from landfill and waste management and 10% from biomass burning.²⁰ Methane has an atmospheric lifetime of about 12 years, so current methane emissions only affect climate for the next few decades. It is thought global methane emissions could be reduced by 25–50% at relatively low cost,²¹ but eliminating the remainder would require behaviour changes whose cost is difficult to determine.²²

including vehicle emissions. It is also a side-effect of methane emissions, increasing their impact on climate by up to 80%.²³ Elevated ozone is hazardous to human and animal health and reduces agricultural crop productivity.

Black carbon aerosols, or soot emissions, are generated from incomplete combustion in sub-standard power stations, vehicle emissions, brick kilns and biomass sources such as wood-fuel cook-stoves. The net climatic impact of many of these processes is unclear: most processes generate mixtures of aerosols, some of which have a warming effect, while others cause cooling.²⁴ Soot and other particulate aerosols represent a significant health hazard: feasible improvements in air quality could prevent between one and five million premature deaths per year, mainly in developing countries.²⁵

Hydrofluorocarbons (HFCs) are the industrial gases introduced to replace the chlorofluorocarbons (CFCs) banned for their impact on the ozone layer. Key HFCs have typical lifetimes of the order of one to 20 years: longer-

lived HFCs are being phased out under the Montreal Protocol. Although their climatic impact to date is limited, HFC emissions are projected to rise substantially. Their role is illustrated below with HFC-134a and HFC-152a.

Table One: Key properties of long-lived and short-lived climate pollutants

Name	Chemical formula	Atmospheric lifetime ²⁶	2011 emissions ²⁷ (Mt = million tonnes)	Impact of past emissions on the planetary energy budget in 2011 ²⁸ (W/m ²)
Carbon dioxide	CO ₂	Centuries to millennia	38,000 Mt CO ₂	1.7
Nitrous oxide	N ₂ O	120 years	7 Mt N ₂ O	0.17
Methane	CH ₄	12 years	330 Mt CH ₄	0.64 (direct)
HFC-134a	CH ₂ FCF ₃	13.4 years	0.04 Mt HFC-134a	<0.01
HFC-152a	CH ₃ CHF ₂	1.5 years	0.03 Mt HFC-152a	Negligible
Tropospheric ozone	O ₃	Weeks	n.a.	0.4 (approx.)
Black carbon	n.a.	Days	2-29 Mt C ²⁹	0.7 (approx.)

1.3 Emission metrics

The climate policy framework established by the UNFCCC allows flexibility in how countries meet their commitments to reduce greenhouse gas emissions. A country whose activities generate large volumes of methane can, for example, opt to reduce those emissions first before reducing CO₂ emissions. Similarly, Emission Trading Systems allow companies to exchange reductions in CO₂ emissions for reductions in other greenhouse gases. In principle, this 'multi-gas basket' approach can identify the most cost-effective measures for reducing greenhouse gas emissions. It requires, however, an exchange rate, known as an emission metric, defining what emissions of different greenhouse gases are worth in terms of each other.³⁰

At present, emissions accounting and trading systems use the 100-year Global Warming Potential, or GWP₁₀₀, to assign relative values to different greenhouse gases. GWP is defined

as the cumulative impact, over a specified time horizon, that the emission of a tonne of a greenhouse gas would have on the planetary energy budget ("how much heat will this tonne trap over a set number of years?"), relative to a tonne of the reference gas CO₂. A tonne of methane emitted today, for example, will have 28 times as much direct impact on the planetary energy budget over the next 100 years as a tonne of CO₂, so methane is assigned³¹ a GWP₁₀₀ of 28. This direct value does not account for the impact of methane on ozone, or the additional CO₂ released by carbon cycle feedbacks, which may be substantial.³²

An alternative metric is the Global Temperature Change Potential, or GTP,³³ which is defined as the impact the emission of a tonne of a greenhouse gas would have on global temperatures at some specified time in the future ("how much warming

will this tonne cause in a given future year?"), again relative to a tonne of CO₂. Since methane has a short lifetime, the impact of a tonne of methane emitted today on temperatures in 100 years' time is small, so the GTP₁₀₀ for methane is seven times lower than its GWP₁₀₀. Both GWP and GTP ignore impacts that occur after the specified time-horizon.

Given that limiting future warming is often cited as the primary goal of climate policy, this discrepancy between GWP and GTP has led some³⁴ to argue that present-day methane emissions may be overvalued by the use of GWP₁₀₀. Over shorter timescales, however, the impact of methane is higher still: a tonne of methane has almost 70 times more impact than a tonne of CO₂ on warming over the ensuing 20 years. If the aim of climate policy were to reduce the global energy imbalance in the short term, then 20-year GWP, GWP₂₀, would be a more appropriate metric. Hence metric choice depends on climate policy priorities.³⁵ It is primarily an issue for short-lived gases: a tonne of nitrous oxide (N₂O), a long-lived gas, is equivalent to about

270 tonnes of CO₂, and this value is relatively independent of metric choice.

Many different metrics have been proposed, reflecting different policy considerations.³⁶ Most of these are found to behave either like GWP or GTP with an appropriate choice of time horizon. In fact, inspection of the table below shows that, for practical purposes, the issue of metric choice is quite simple: for all gases shown except for the very short-lived HFC-152a, values of GWP₁₀₀ are similar to their 40-year GTPs. Hence GWP₁₀₀ may be understood, informally, as a measure of the relative impact of different greenhouse gases on temperature rise over the next 40 years or less, while GTP₁₀₀ is a measure of relative impact on temperatures in the early 22nd century. The choice between GWP₁₀₀ and GTP₁₀₀ is therefore, to a good approximation, simply a decision as to whether we wish to prioritise reducing climate change over the next few decades or reducing climate change in the more distant future.

Table Two: 'Greenhouse gas exchange rates', or what a tonne of each gas is worth in terms of tonnes of CO₂ under various climate metrics, for the three most important greenhouse gases and two illustrative HFCs.³⁷

Gas	Global Warming Potential		Global Temperature Change Potential		
	GWP ₂₀	GWP ₁₀₀	GTP ₂₀	GTP ₄₀	GTP ₁₀₀
Carbon dioxide	1	1	1	1	1
Nitrous oxide	264	265	277	285	234
Methane	84	28	67	26	4
HFC-134a	3710	1300	3050	1173	201
HFC-152a	506	138	174	36	19
Black carbon ³⁸	3200	910	925	n.a.	130

2. The different impacts of SLCP and CO₂ mitigation on global temperatures

This section explains the potential impact of immediate reductions in SLCP and CO₂ emissions on future climate change. Figure One shows three scenarios for CO₂ emissions (based on the Representative Concentration Pathways³⁹ – RCPs – used by the IPCC). Under the high emissions scenario, in red, global CO₂ emissions continue to increase at about 1.8–2% per year. The solid blue line represents an ambitious mitigation scenario under which climate policies are introduced to meet the goal of avoiding more than 2°C of warming relative to pre-industrial temperatures. The dashed line shows a “delayed mitigation scenario” which assumes the same percent-per-year reductions in emissions as the solid blue line, but starting 20 years later,⁴⁰ in 2035.




Figure One: Idealised CO₂ emission scenarios used in this policy paper. Red scenario based on IPCC RCP8.5 scenario. Solid blue scenario assumes same percent-per-year rates of change of emissions as projected in RCP3PD from 2010, but starting in 2015. Dashed scenario computed likewise, but starting in 2035.

The solid lines in Figure Two show the potential impact on global temperatures⁴¹ of the solid red and solid blue scenarios in Figure One assuming the same scenario for non-CO₂ climate drivers in both cases. This paper uses the average of a ‘high

(RCP8.5) and an ‘ambitious mitigation’ (RCP3PD) scenario for these non-CO₂ drivers, which gives approximately stable methane and nitrous oxide emissions from 2020 onwards. It is, of course, artificial to assume strenuous efforts to reduce CO₂ have no impact on efforts to reduce other emissions, but this is an idealised example to illustrate the respective impact of CO₂ and non-CO₂ mitigation.⁴²




Figure Two: Temperature response to idealised CO₂ emission profiles with and without immediate reductions in short-lived climate pollutants. Red implies high CO₂ emissions scenario (solid red line in Figure One), blue implies ambitious CO₂ mitigation scenario (solid blue line in Figure Two). Solid lines assume approximately stable SLCP emissions from 2020 onwards. Dashed lines assume ambitious measures to reduce methane and black carbon emissions over the period 2015–2035. Dashed blue line assumes both ambitious CO₂ and SLCP measures occur. Dashed purple line is the same as the dashed blue scenario except that methane reductions are offset by an “equivalent” increase in CO₂ emissions using the GWP₁₀₀ metric.

The solid red scenario shows temperatures continuing to rise rapidly throughout the 21st century and beyond. Under the solid

blue scenario, global temperatures stabilise by the end of the century as CO₂ emissions approach zero, but the goal of keeping global temperatures below 2°C is missed (under this particular model of the response).

The dashed red and blue lines show warming under the same scenarios for CO₂ emissions but, instead of increasing, methane emissions are reduced to 75% of their 2015 values over the 20 years 2015–2035 and held constant thereafter, and measures to reduce black carbon emissions result in a further 0.25 W/m² reduction in non-CO₂ climate warming over the same period.⁴³ This corresponds to a relatively optimistic scenario for reducing SLCP emissions, although comparable to published estimates of what might be possible, just as the solid blue scenario in Figure One is at the optimistic end of scenarios for reducing CO₂ emissions. Detailed assessment of the cost of these illustrative scenarios is beyond the scope of this paper, but it should be noted that estimates of the cost of SLCP reductions of this magnitude are substantially lower than estimates of the cost of these CO₂ reductions.

Figures such as this one have been used to justify the claim that, as the United Nations Environment Programme/World Meteorological Organization report⁴⁴ put it, “near-term emission control measures [on methane and soot], together with measures to reduce CO₂ emissions, would greatly improve the chances of keeping Earth’s temperature increase to less than 2°C.” Even with very ambitious mitigation, reductions in CO₂ emissions take some decades to impact global temperatures because CO₂ emissions accumulate in the climate system. Reductions in SLCP emissions have a more immediate impact, so the red dashed line (with SLCP measures, but no CO₂ mitigation) remains below the blue solid line (only CO₂ measures) beyond 2050.

Hence if the sole focus of climate policy were to limit warming to 2050, then these SLCP measures would undoubtedly be far more cost-effective than these CO₂ reductions. If we

consider other proposed climate targets such as global energy imbalance⁴⁵ in 2050, a similar picture emerges: these SLCP measures would reduce the global energy imbalance by a similar amount (0.8 W/m²) to these CO₂ emission reductions, at a much lower cost. It is clear from the figure, however, that a world in which the global energy imbalance has been reduced by 0.8 W/m² through SLCP measures would be in a very different position in 2050 to a world in which the same reduction had been achieved with CO₂ emission reductions: the former would be still warming rapidly, while the latter would be approaching climate stabilisation. Hence setting goals for temperatures or energy imbalance in 2050 will never be sufficient if policies are also concerned with what happens thereafter.

A danger with idealised modelling studies such as that shown in Figures Two (above) and Three (below) is that they conceal the importance of natural, and unpredictable, climate variability. Recent studies⁴⁶ have shown that the impact of such variability can be high, particularly on climate changes on regional scales, potentially overwhelming any externally driven trend on 20- to 40-year timescales even under a relatively high forcing scenario. Hence the climatic benefits of SLCP mitigation on these timescales should be kept in perspective: SLCP mitigation could delay the time at which a threshold is crossed by up to 20 years, but internal climate variability could also delay, or bring forward, that time by a similar margin.

The combination of aggressive CO₂ and SLCP mitigation (dashed blue line) remains comfortably below the 2°C ‘guardrail’, but it must be stressed that this assumes both sets of mitigation measures are taken, and that no resources are diverted from the CO₂ mitigation effort by SLCP mitigation. The dashed purple line shows what happens if SLCP mitigation measures are taken as in the dashed blue line, but those achieving these rapid methane reductions are allowed to ‘sell’ or offset them against CO₂ reductions, using the conventional GWP₁₀₀ metric of exchange.⁴⁷ There would be

a clear incentive for this to occur unless it were explicitly prohibited, since methane emissions are already routinely traded against CO₂ emissions using the GWP₁₀₀ metric, and under an aggressive mitigation scenario such as this one, the incentives to find alternatives to CO₂ mitigation would be considerable. Such large-scale offsetting, if it were to occur, would wipe out the impact of methane reductions in the short term, while in the longer term it would result in substantially more warming because the additional CO₂ released continues to warm the climate long after the impact of avoided methane emissions has dissipated.

In the absence of CO₂ mitigation, shown by the dashed red line, SLCP measures only delay, but do not prevent, temperatures crossing any particular threshold. The red dashed line tracks the red solid line (no mitigation effort at all) 15–20 years later. This has prompted some, in the popular press, to suggest that action on SLCPs could “give politicians two extra decades to tackle the less tractable question about what to do about CO₂”,⁴⁸ but this is a serious misinterpretation, as this paper will show in the next section.

3. The impacts of delaying emission reductions

Figure Three compares the impact of the CO₂ and SLCP emission measures shown in Figure Two with the impact of the same measures initiated 20 years later.⁴⁹ The negative impact of a 20-year delay in CO₂ mitigation is much greater than the positive impact of any possible SLCP measures that might be taken in the meantime. Since CO₂ accumulates in the climate system, as long as emissions continue to increase at 1.8–2%/year as they are doing at present, the eventual peak warming to which we are committed is increasing at the same rate, which is much faster than observed temperatures are increasing.⁵⁰

The CO₂ emission reduction rates that would have met the goal of limiting warming to 2 °C if initiated in 2010 would only meet a 2.2 °C goal if initiated in 2015, when emissions will be 10% higher. If CO₂ emissions continue to rise as they are doing, a 20-year delay would add almost 50% to eventual peak warming, shown by the difference between the red scenarios (delayed CO₂ measures) and blue scenarios (early CO₂ measures) in Figure Three. The only exception to this rule would be if a period of delay were used explicitly to invest in technologies that allowed CO₂ emissions to be reduced much more rapidly after the emissions peak (accepting the additional risk that such rapid reductions might not prove possible).




Figure Three: Impact of delayed emission cuts on global temperatures. Dashed blue line: CO₂ and SLCP measures initiated immediately, in 2015, as in Figure Two. Solid blue line: SLCP cuts delayed to 2035, CO₂ cuts initiated in 2015. Dashed red line: CO₂ cuts delayed to 2035, SLCP cuts initiated in 2015. Solid red line: both delayed to 2035.

The impact of delaying SLCP mitigation measures is very different. As would be expected, delaying SLCP measures by 20 years results in more warming over the 2015–2035 period: the difference between the solid lines and the dashed lines. In the longer term, however, the impact of the timing of SLCP measures on peak warming depends on what happens to CO₂ emissions in the meantime. If CO₂ emissions are reduced immediately (blue lines), then failing to reduce SLCP emissions at the same time could add a few tenths of a degree to peak warming (blue solid versus blue dashed lines). But if CO₂ emission reductions are also delayed for 20 years, it makes no difference to peak warming whether SLCP measures are implemented immediately or deferred until after 2035 (red solid versus red dashed lines).

Again, this is a simple illustration of a more general point: SLCP emissions only have an impact on peak warming under circumstances

in which CO₂ emissions are either already falling or about to fall rapidly.⁵¹ Hence, if the main objective of climate policy is to limit peak warming, then SLCP mitigation could be delayed until after global CO₂ emissions have started to decline and are expected to continue to decline towards zero. It is quite wrong to suggest that bringing forward SLCP emission reductions can “buy time” to procrastinate over CO₂: quite the reverse. Early SLCP measures only have an impact on peak warming if ambitious CO₂ mitigation measures are already under way.

4. The implications for climate metrics

The simple scenarios shown in Figures One, Two and Three suggest that ambitious measures to reduce SLCP emissions like methane over the next 20 years would have more of an impact on global temperatures out to 2050 than equally ambitious (and potentially much more expensive) cuts in CO₂ emissions. On the other hand, the scenarios also show that unless CO₂ emissions are being reduced at the same time, it makes almost no difference to peak warming whether SLCP emissions are reduced now, or reduced in some later decade closer to the time that temperatures peak. So what is a tonne of methane emitted today ‘worth’ relative to a tonne of CO₂?

The example shows that there can be no exact equivalence between any given amount of methane emissions and a given amount of CO₂. Any decision on a metric or ‘exchange rate’ stating what a tonne of methane emissions is worth in terms of equivalent CO₂ depends not only on the properties of the two gases (their atmospheric lifetimes and relative potency as greenhouse gases), but also on assumptions about how the climate system will respond to emissions, and decisions about the relative importance of impacts on different time-scales.

This much is well known. What is less widely appreciated is that the appropriate metric to use today also depends on future emissions. If policies like the goal of avoiding more than 2 °C of warming refer to maximum climate change independent of timescale, then assumptions about future emissions, which determine the timing of peak warming, affect priorities today. Under these conditions, the choice of metric represents, at some level, a bet on the success or failure of future climate mitigation policy. Hence there is no purely technical method of determining the correct metric of exchange between a tonne of methane and a tonne of CO₂. Whatever metric is adopted may have distortionary and potentially surprising implications for mitigation outcomes.⁵²

One option that has been proposed⁵³ that would better reflect the actual climatic impact of CO₂ and SLCPs would be to allow a tonne of CO₂ to be released in exchange for a *permanent* reduction in the *rate* of emission of a SLCP. It is difficult to see how this could be implemented in an emission-trading scheme because, while emissions in any given year can be verified, it is unclear how an indefinite commitment to reduce an emission rate could be enforceable. Nevertheless, in comparing policies, it may be useful to recall that every tonne of CO₂ released has an effectively permanent impact on climate, while only permanent changes in SLCP emission rates have a comparable long-term impact.⁵⁴

5. The impact of metrics on mitigation priorities: are SLCPs like methane over-valued or under-valued?

The implications of metric choice on mitigation priorities are illustrated in Figure Four, which shows 2011 emissions of these three main greenhouse gases expressed as CO₂-equivalent emissions using GWP₂₀, GWP₁₀₀ and GTP₁₀₀ respectively. Under the GWP₂₀ metric, 2011 methane emissions are equivalent to almost 30 billion tonnes of CO₂ per year, over two thirds of 2011 CO₂ emissions. The current standard metric, GWP₁₀₀, assigns three times lower weight to methane emissions than GWP₂₀, while the GTP₁₀₀ metric assigns a weight to current methane emissions that is seven times lower still.




Figure Four: Emissions of three main greenhouse gases in 2011 expressed in terms of CO₂-equivalent using the GWP₂₀, GWP₁₀₀ and GTP₁₀₀ metrics.

Variation of the nominal impact of current methane emissions by a factor of 20 makes the issue of metric choice seem impossibly arbitrary, but in fact these numbers can be easily understood by reference to the example in Figure Two. GWP₂₀ is a measure of the impact of greenhouse gas emissions on energy imbalance and warming rates over the coming 20 years, so the high value of

methane emissions under GWP₂₀ reflects the observation in Figure Two that a halving of methane emissions would be as effective in reducing warming over the following couple of decades as an immediate 40% reduction in CO₂ emissions. Hence if the sole priority of climate policy is to reduce warming or global energy imbalance over the next 20 years, it could be argued that GWP₁₀₀ under-values current methane emissions, and we should instead use a metric like GWP₂₀.

In contrast, if the aim of climate policy is to limit warming in 100 years' time, then the relevant metric is GTP₁₀₀, which suggests that a 50% reduction in 2011 methane emissions would be equivalent to only a 2.5% reduction in CO₂ emissions. Again, this can be understood by the example in Figure Three: it makes almost no difference to temperatures at the end of the century whether SLCP measures are taken now or taken in 20 years' time: a tonne of methane emitted today has almost no impact on temperatures in 2100. Hence if 100-year warming is the main focus of climate policy, then GWP₁₀₀ vastly overstates the importance of current methane emissions.

Hence a case can be made that the GWP₁₀₀ overstates the importance of present-day methane emissions, and also that it understates it, depending on whether the aim of climate policy is to limit warming in 2100 or to limit climate change over the next 20 years. This is essentially restating, in terms of climate metrics, the results shown in Figures Two and Three: the relative importance of present-day SLCP emissions depends on the time-frame of interest, and if the goal is to limit peak warming, then it also depends on the success of future mitigation policies that determine when peak temperatures are reached.

6. The special case of methane, GWP₁₀₀ and the 2 °C goal

GWP₁₀₀, despite its name, is an indicator of the relative impact of present-day methane and CO₂ emissions on temperature rise between now and the 2050s: the reason is that methane's GWP₁₀₀ is close to the value of its GTP₄₀, which is a measure of the relative impact of current emissions on temperatures roughly 40 years from now.⁵⁵

The increase in temperatures between now and the 2050s matters for the goal set by the UNFCCC in Cancun in 2010 of limiting human-induced warming to 2 °C above pre-industrial, because temperatures are already about 0.8 °C above pre-industrial and are projected to rise by about another half a degree by 2030.⁵⁶ If the 2 °C goal is to be met, then temperatures will have to start to stabilise by mid-century. The impact of emissions today on temperatures in the 2050s is thus a very approximate indicator of their impact on peak warming, *if and only if* emissions of both CO₂ and other climate pollutants are reduced substantially over the coming decades to stabilise temperatures shortly thereafter.

The argument can be made, therefore, that GWP₁₀₀ represents a reasonable metric to use to compare the impact of different gases on peak warming, but *only* under stringent mitigation scenarios that achieve the aspirational 2 °C goal now adopted by the Parties to the UNFCCC. Use of the GWP₁₀₀ metric (which implies a relatively high priority to SLCP mitigation over the coming decade) therefore *assumes* that the 2 °C goal will in fact be met. If CO₂ emissions are not reduced in the near term and temperatures continue to rise through 2070 and beyond, the impact of present-day methane emissions on peak warming is vastly overstated by GWP₁₀₀.

As long as aggregate national commitments to reduce greenhouse gas emissions are insufficient to meet the 2 °C goal, as is the

case at present,⁵⁷ continued use of GWP₁₀₀ as a metric to compare and exchange emission reductions of different gases implies that the true goal of climate policy is to minimise the increase in temperatures over the next 40 years or so, and not to limit peak warming at all. This would be a coherent policy objective, but if it has become the primary objective of governments engaged in the UNFCCC process, they should arguably say so. It is very misleading to base climate policy on the assumption that the 2 °C goal will be met in the absence of any concrete plan for meeting it.

7. The implications of metric choice for carbon markets

Much of the interest in SLCP emissions arises from the fact that these emissions could be reduced at relatively low cost compared to the equivalent quantity of CO₂ emissions. These 'Marginal Abatement Costs'⁵⁸ are very uncertain, but the cost differentials are very thought to be so great that many SLCP emission reductions would still appear desirable whatever metric is used to compare them to CO₂ emissions.⁵⁹ If, however, SLCP emissions are traded or offset against CO₂ while climate policy goals are still set in terms of total CO₂-equivalent emissions, as in the purple dashed line in Figure Two, then metric choice could matter a great deal through its impact on CO₂ emissions, as this section explains.

Suppose, for the sake of an idealised example, that 50% of current methane emissions can be eliminated at very low cost while the remaining 50% would be much more difficult and expensive to eliminate. Suppose also that nitrous oxide emissions are prohibitively expensive to reduce.

Figure Five shows the implications for CO₂ emissions of a 10% and 50% reduction in total CO₂-equivalent emissions of these three gases, using three different metrics to calculate CO₂-equivalence. Under GTP₁₀₀, CO₂-equivalent emissions are almost entirely dominated by CO₂ itself, so any reduction in CO₂-equivalent emissions requires reductions in CO₂. Under GWP₁₀₀, a 10% reduction in total CO₂-equivalent emissions could be achieved with methane alone, leaving actual CO₂ emissions almost unchanged. Under GWP₂₀, a 10% reduction in total CO₂-equivalent emissions could be accompanied by a 20% increase in CO₂ emissions.




Figure Five: Impacts of 10% and 50% cuts in 2011 total 'CO₂-equivalent' emissions under different emission metrics. It is assumed that 50% of methane emissions can be abated at very low cost, and that the remainder, plus nitrous oxide emissions, are very expensive to abate. CO₂ emissions are calculated to give correct total emission cuts, expressed in terms of CO₂-equivalence using the GWP₂₀, GWP₁₀₀ and GTP₁₀₀ metrics.

For a 50% reduction in CO₂-equivalent emissions, CO₂ emissions themselves have to be reduced no matter what metric is used to define CO₂-equivalence. Hence, in the long term, meeting climate policy goals will require very substantial reductions in all greenhouse gas emissions regardless of the metric used to compare them.⁶⁰ In the short term, however, the greater the 'value' assigned to methane emissions, the less CO₂ mitigation will be required for any given reduction in total CO₂-equivalent emissions: the impact of metric choice on methane emissions is not whether or not methane emission reductions occur, but the volume of CO₂ emissions that countries and companies are allowed to offset against them.

8. The implications of metric choice for the timing of emission reductions

Figure Six shows how metric choice might affect the timing of mitigation decisions under an ambitious mitigation scenario. In this example, total CO₂-equivalent emissions are required to peak within eight years of 2015 and then decline steadily to less than half 2015 emissions by 2050. Methane emissions are reduced by 50% over 2015–2035, nitrous oxide emissions are unchanged and CO₂ emissions adjusted to make up the remainder.





Figure Six: Emissions of the three main greenhouse gases under an idealised ambitious mitigation scenario defined in terms of total CO₂-equivalent emissions but using three different metrics to define CO₂-equivalence.

When CO₂-equivalence is defined in terms of either GWP₁₀₀ or GTP₁₀₀ (panels a and b), this yields similar reductions in all three gases, at least out to 2050. So, under an ambitious mitigation scenario, the same decisions would be made at a global level to meet a given total CO₂-equivalent emission target whether GWP₁₀₀ or GTP₁₀₀ is used, although the choice might still have substantial implications for how the burden of reductions is distributed across countries and sectors.

When CO₂-equivalence is defined in terms of GWP₂₀ (panel c), corresponding to a policy focused on short-term goals for global energy imbalance rather than temperature,⁶¹ this yields a rather different scenario. Methane emissions are still reduced by 50% by 2035, but because they play a much larger nominal role in present-day emissions, the target for total CO₂-equivalent emissions can be met while allowing CO₂ emissions themselves to continue to rise. At that point, because of the assumed high cost of further methane reductions, CO₂ emissions have to fall sharply.

In the short term, using a metric that assigns a relatively high weight to methane and other SLCPs gives the impression that climate mitigation can be achieved relatively cheaply. This might be politically advantageous in terms of building momentum for climate mitigation,⁶² and might not matter in the long term *provided* CO₂ emissions are successfully and rapidly reduced when they need to be.

Any delay in initiating CO₂ emission reductions represents, however, a substantial risk, because we still do not know how difficult and costly these reductions will be, either technically or politically. Long-term climate change is overwhelmingly determined by cumulative CO₂ emissions, so the longer actual reductions in CO₂ emissions are postponed, the more difficult it becomes to limit long-term warming. The same rate of CO₂ emission reductions that would limit CO₂-induced warming to 3 °C if initiated now would only limit it to 4 °C if initiated after 15 more years of emissions growth at 2% per year.

9. Conclusion and recommendations

The authors of the UNFCCC recognised that the goals of climate policy could not be summed up in a single sentence. Article 2 of the Convention states:

“The ultimate objective of this Convention [is] stabilization of greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system. Such a level should be achieved within a time-frame sufficient to allow ecosystems to adapt naturally to climate change, to ensure that food production is not threatened and to enable economic development to proceed in a sustainable manner.”

The first sentence was, and remains, a commitment to future generations. The greatest risks of dangerous anthropogenic interference in the climate system are unlikely to manifest themselves within the lifetime of anyone who was alive when the Convention was opened for signature in 1992, and very possibly not in the lifetime of anyone alive today in 2015. But the second sentence recognises more immediate concerns: allowing ecosystems to adapt and ensuring food production and economic development can continue.

Reduced to its simplest form, the debate over emission metrics and SLCP versus CO₂ mitigation can be conceived as addressing the two sentences of this Article. It is necessary to limit cumulative emissions of CO₂ to stabilise climate and hence limit the risk of dangerous anthropogenic interference in the climate system in the long term. But reducing SLCP emissions may well be a more cost-effective way to limit the rate of climate change over the coming decades to ensure that ecosystems, food production and the economy can adapt, which also has a role in avoiding dangerous climate change.

Proponents of early action on SLCP emissions frequently emphasise the ‘complementary’

nature of SLCP and CO₂ mitigation, but it is important to be clear what this complementarity means: they are not two ways of achieving the same goal, but address fundamentally different goals, affecting different generations. Early action on SLCP mitigation could affect the temperatures and climate impacts experienced by the generation of today’s decision-makers, but will have little impact on the warming experienced by future generations. Unless it is accompanied by ambitious reductions in CO₂ emissions, early SLCP mitigation will also have very little impact on eventual peak warming.

Just as the objective of the UNFCCC could not be summed up in a single sentence, so there is no single emission metric that makes reducing emissions of SLCPs like methane equivalent to reducing emissions of CO₂. They have different objectives, emphasising the interests of different generations, and there is no alternative to designing and monitoring climate policies to ensure that *both* of these vital objectives are met.

To meet the goals of the UNFCCC, policies are required to ensure that global CO₂ emissions are contained within a cumulative budget consistent with limiting warming to a safe level. These policies must be independent of, and in addition to, any multi-gas emission goals. In effect, this implies a ‘peak CO₂ first’ strategy: the need to limit cumulative CO₂ emissions would over-ride most opportunities to offset CO₂ reductions against SLCP measures until global CO₂ emissions are falling fast enough that there is a realistic prospect of meeting the cumulative budget. As soon as those conditions are met (for example, when CO₂ emissions are projected to reach zero before global temperatures reach 2 °C), SLCP emission reductions will become a crucial priority to limit peak warming.

Notes

- 1 Stocker, T.F., Qin, D. *et al* (2013): *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge and New York, NY.
- 2 Myhre, G. *et al* (2013): Anthropogenic and natural radiative forcing. In Stocker, T.F., Qin, D. *et al*. *op. cit.* Estimates suggesting slower very recent rates of increase (<http://www.bbc.co.uk/news/science-environment-31872460>) remain highly provisional.
- 3 Victor, D. G., Kennel, C. F. & Ramanathan, V. (2012): The climate threat we can beat. *Foreign Aff.*, 91, 112–114.
- 4 Shindell, D. *et al* (2012): Simultaneously mitigating near-term climate change and improving human health and food security, *Science*, 335, 183–189.
- 5 Schmale, J. *et al* (2014): Air pollution: Clean up our skies, *Nature*, 515, 335–337.
- 6 Victor, D. G. & Kennel, C. F. (2014): Climate policy: Ditch the 2°C warming goal, *Nature*, 514, 30–31.
- 7 UNEP Climate and Clean Air Coalition (2014): *Time to Act to Reduce Short Lived Climate Pollutants*, http://www.unep.org/ccac/Portals/50162/docs/publications/Time_To_Act/SLCP_TimeToAct_lores.pdf
- 8 See, for example, Government of Mexico (2015): *Intended Nationally Determined Contribution, Submission to the UNFCCC*, <http://www4.unfccc.int/submissions/INDC/Published%20Documents/Mexico/1/MEXICO%20INDC%2003.30.2015.pdf>
- 9 Collins, M., Knutti, R. *et al* (2013): Long-term climate change: projections, commitments and irreversibility. In Stocker, T. F., Qin, D. *et al*, *op. cit.*
- 10 Allen, M. R. *et al* (2009): Warming caused by cumulative carbon emissions towards the trillionth tonne, *Nature*, 458, 1163–1166.
- 11 Rogelj, J. *et al* (2014): Disentangling the effects of CO₂ and short-lived climate forcer mitigation, *Proc. Nat. Acad. Sci.*, 111, 16325–16330.
- 12 Shoemaker, J. K. & Schrag, D. P. (2013): The danger of overvaluing methane's influence on future climate change, *Clim. Change*, 120, 903–914.
- 13 Pierrehumbert, R. T. (2014): Short-lived climate pollution, *Ann. Rev. Earth & Planetary Sciences*, 42, 341–379. This comprehensive paper provides a more technical analysis of many of the issues discussed in this report.
- 14 Ciais, P., Sabine, C. *et al* (2013): Carbon and other biogeochemical cycles. In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 15 Ciais, P., Sabine, C. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 16 Collins, M., Knutti, R. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 17 Clarke, L., Jiang, K. *et al* (2014): Assessing transformation pathways. In *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Edenhofer, O., Pichs-Madruga, R. *et al* (eds.)]. Cambridge University Press, Cambridge and New York, NY.
- 18 Ciais, P., Sabine, C. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.* All lifetimes quoted here to two significant figures only.
- 19 Clarke, L., Jiang, K. *et al* (2014). In Edenhofer, O., Pichs-Madruga, R. *et al*, *op. cit.*
- 20 Ciais, P., Sabine, C. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 21 Shindell, D. *et al* (2012), *op. cit.*
- 22 Myhre, G. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 23 Myhre, G. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 24 Bond, T. C. *et al* (2013): Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res. Atmos.*, 118, 5380–5552.
- 25 Shindell, D. *et al* (2012), *op. cit.*
- 26 Ciais, P., Sabine, C. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 27 IPCC (2014): *Climate Change 2014: Synthesis Report* [Pachauri, R. K. & Meyer, L. (eds.)]. Geneva, p. 151.
- 28 Myhre, G. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 29 Emission figures for black carbon from Bond, T. C. *et al* (2013), *op. cit.*
- 30 Fuglestedt, J. *et al* (2003): Metrics of climate change: Assessing radiative forcing and emission indices, *Clim. Change*, 58, 267–331.
- 31 Myhre, G. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 32 Gillett, N. & Matthews, H. (2010): Accounting for carbon cycle feedbacks in a comparison of the global warming effects of greenhouse gases, *Environ. Res. Lett.*, 5, 034011.
- 33 Shine, K., Fuglestedt, J., Hailemariam, K. & Stuber, N. (2005): Alternatives to the global warming potential for comparing climate impacts of emissions of greenhouse gases, *Clim. Change*, 68, 281–302.
- 34 Shoemaker, J. K. & Schrag, D. P. (2013) *op. cit.*
- 35 See box 3.2 in Pachauri, R. K. & Meyer, L. (eds.) (2014), *op. cit.*, pp. 87–88.
- 36 Tol, R. S. J. *et al* (2012): A unifying framework for metrics for aggregating the climate effect of different emissions, *Environ. Res. Lett.*, 7, 044006.
- 37 Metric values except for black carbon from Myhre, G. *et al* (2013), *op. cit.*, with GWP₁₀₀ computed using the algorithm provided in the Supplementary Online Material.
- 38 Metric values for black carbon as given in Bond, T. C. *et al* (2013), *op. cit.* Note that these exclude the cooling climatic impact of other pollutants generated by many of the activities that generate black carbon emissions and hence may overstate the apparent warming impact of these activities.
- 39 Van Vuuren, D. P. *et al* (2011): The representative concentration pathways: An overview, *Clim. Change*, 109, 5–31. The standard RCP3PD scenario envisages emission reductions beginning in 2010, whereas in fact emissions have continued to track the RCP8.5 scenario since then, so this paper has assumed the same percent-per-year changes in emissions that are assumed in RCP3PD, but delayed by five years, starting in 2015.
- 40 Allen, M. R. & Stocker, T. F. (2013): Impact of delay in reducing carbon dioxide emissions, *Nature Climate Change*, 4, 23–26.
- 41 Using the simple impulse response model of Boucher, O. & Reddy, M. (2008): Climate trade-off between black carbon and carbon dioxide emissions, *Energy Policy*, 36, 193–200, with an equilibrium climate sensitivity of 3°C and a Transient Climate Response of 1.8°C, corresponding to the mean response of more complex models in the CMIP5 ensemble.
- 42 In the time-series examples in this policy paper, emissions are inferred from the RCP radiative forcing profiles using the lifetime and efficacy factors in Myhre, G. *et al* (2013), *op. cit.* Although adequate for a heuristic example, assuming constant lifetimes for all gases neglects important feedbacks, such as the feedback between rising temperatures and the release of CO₂ from natural storage. Emissions inferred in this way understate past emissions and overstate future emissions relative to more complex models but give the correct impact on the planetary energy budget. Figures Four and Five use actual reported emissions data.
- 43 This corresponds approximately to the “Early SLCP measures” scenario in Shindell, D. *et al* (2013), *op. cit.* The IPCC's RCPs do not include a specific scenario of rapid early reduction of SLCP emissions. Net non-CO₂ climate forcing remains around present-day levels until after 2050 even under the RCP3PD mitigation scenario.
- 44 UNEP & WMO (2011): *Integrated Assessment of Black Carbon and Tropospheric Ozone*, UNEP, Nairobi.
- 45 Victor, D. G. & Kennel, C. F. (2014), *op. cit.*
- 46 Deser, C., Phillips, A. S., Alexander, M. A. & Smoliak, B. V. (2014): Projecting North American climate over the next 50 years: Uncertainty due to internal variability, *J. Climate*, 27, 2271–2296.
- 47 Pierrehumbert, R. T. (2014), *op. cit.*
- 48 Global warming: The new black, *The Economist* (19 January 2013).
- 49 Bowerman, N. H. A. *et al* (2013): The role of short-lived climate pollutants in meeting temperature goals, *Nature Climate Change*, 3, 1021–1024.
- 50 Allen, M. R. & Stocker, T. F. (2013), *op. cit.*
- 51 Bowerman, N. H. A. *et al* (2013), Pierrehumbert, R. T. (2014), and Allen, M. R. & Stocker, T. F. (2013), *op. cit.*
- 52 Daniel, J. S. *et al* (2012): Limitations of single-basket trading: Lessons from the Montreal Protocol for climate policy, *Clim. Change*, 111, 241–248.
- 53 Smith, S. M. *et al* (2012): Equivalence of greenhouse-gas emissions for peak temperature limits, *Nature Climate Change*, 2, 535–538.
- 54 Pierrehumbert, R. T. (2014), *op. cit.*
- 55 See figure 8.30 of Myhre, G. *et al* (2013). In Stocker, T.F., Qin, D. *et al*, *op. cit.*
- 56 See figure TS.14 in Stocker, T.F., Qin, D. *et al* (2013), *op. cit.*
- 57 UNEP (2014): *The Emissions Gap Report 2014*, UNEP, Nairobi <http://www.unep.org/emissionsgapreport2014/>
- 58 Estimates of the cost of reducing emissions, as a function of amount of reduction, or Marginal Abatement Costs, are highly uncertain: the figures of 50% relatively low-cost methane abatement and inflexible nitrous oxide emissions are used for illustration, based on the UK Department of Energy and Climate Change GLOCAF model derived from the IMAGE model of van Vuuren, D. P. *et al* (2004): Responses to technology and taxes in a simulated world, *Energy Economics*, 26(4), 579–601.
- 59 Huntingford, C. *et al* (2015): The implications of carbon dioxide and methane exchange for the heavy mitigation RCP2.6 scenario under two metrics, *Environmental Science and Policy*, 51, 77–87.
- 60 Reisinger, A. R. *et al* (2013): Implications of alternative metrics for global mitigation costs and greenhouse gas emissions from agriculture, *Clim. Change*, 117, 677–690.
- 61 Victor, D. G. & Kennel, C. F. (2014), *op. cit.*
- 62 Victor, D. G., Kennel, C. F. & Ramanathan, V. (2012), *op. cit.*

About the author

Professor Myles Allen is Co-Director of the Oxford Martin Programme on Resource Stewardship. He is Professor of Geosystem Science in the School of Geography and the Environment and the Department of Physics at the University of Oxford, Leader of the Environmental Change Institute's Climate Research Programme and a Fellow of Linacre College. His research focuses on how human and natural influences on climate contribute to observed climate change and risks of extreme weather, and quantifying their implications for long-range climate forecasts.

Myles Allen has served on the Intergovernmental Panel on Climate Change as Lead Author on *Detection of Climate Change and Attribution of Causes* for the 3rd and 5th Assessments in 2001 and 2013, as Review Editor on *Global Climate Projections* for the 4th Assessment in 2007 and as a member of the Synthesis Report Core Writing Team in 2014. He proposed the use of Probabilistic Event Attribution to quantify the contribution of human and other external influences on climate to specific individual weather events and leads the www.climateprediction.net project, using distributed computing to run the world's largest ensemble climate modelling experiments.

Acknowledgements

The author would like to express his gratitude to the reviewers of this paper, Professor Raymond Pierrehumbert, Dr Andy Reisinger and Professor David Victor. This work benefited from discussions with and suggestions from Djamila Amimer, Niel Bowerman, Bill Collins, Piers Forster, Jan Fuglestedt, Laila Gohar, Nick Howarth, Chris Huntingford, David Lee, Jason Lowe, Alex Otto and Keith Shine, through research undertaken for the UK Department of Energy and Climate Change under contract TRN 307/11/2011 "Assessing the Options for Greenhouse Gas Metrics", but the views expressed herein are the sole responsibility of the author. He also thanks the members of the Oxford Martin Policy and Research Steering Group, and Anushya Devendra for editorial and logistical support.

Agreements and strategies addressing climate change involve setting priorities between reducing emissions of different climate pollutants. While avoiding dangerous climate change ultimately requires reducing emissions of long-lived greenhouse gases like carbon dioxide to zero, action on short-lived climate pollutants such as methane and soot may be a cheaper way of mitigating warming over the next few decades, with substantial co-benefits to agriculture and human health. This paper explains how the 'short-lived' versus 'long-lived' discussion is not really a technical issue at all, but an expression of inter-generational priorities; the standard policy framework or metric used to compare different climate pollutants, the '100-year Global Warming Potential', prioritises the interests of the current generation over future generations. The paper provides a non-technical overview and sets out policy recommendations, notably a 'peak CO₂ first' strategy, as countries and companies prepare for the UN Climate Change Conference in Paris in December 2015.

Oxford Martin Policy Papers harness the interdisciplinary research of the Oxford Martin School to address today's critical policy gaps. The development of these policy papers is overseen by the Oxford Martin Policy and Research Steering Group, chaired by Professor Sir John Beddington FRS, Senior Adviser to the Oxford Martin School. This paper reflects the views of the author, and does not necessarily reflect the position of this Steering Group, the Oxford Martin School or the University of Oxford.



Oxford Martin School
University of Oxford
Old Indian Institute
34 Broad Street
Oxford OX1 3BD
United Kingdom

Tel: +44 (0)1865 287430
Fax: +44 (0)1865 287435
Email: info@oxfordmartin.ox.ac.uk
Web: <http://www.oxfordmartin.ox.ac.uk/>
Twitter: @oxmartinschool
Facebook: <http://www.facebook.com/oxfordmartinschool>

© Oxford Martin School,
University of Oxford, 2015

This paper was designed and printed
by Technique Print Group.



NonCommercial-NoDerivs 3.0 Unported Licence.
To view a copy of this licence, visit
<http://creativecommons.org/licenses/by-nc-nd/3.0/>